



 RIDLEY

BARASTOC

ESSENTIAL FEEDS

**CHICK
STARTER**

COMPLETE & BALANCED
CRUMBLE



WHAT WILL IT PROVIDE MY CHICKS?

Nutritionally balanced starter feed specifically formulated to support the growth and health of chicks during the first few weeks post hatching.

Contains an anticoccidial as an aid in the prevention of coccidiosis, an intestinal infection caused by the Eimeria parasite which young birds are vulnerable to.


Presented as a small crumble which is suitable for small beaks.

BARASTOC

ESSENTIAL FEEDS

CHICK STARTER

COMPLETE & BALANCED CRUMBLE



NUTRIENT LEVELS

TYPICAL ANALYSIS AS FED PER KG

NUTRIENT LEVELS			
Min Crude Protein	19.5%	Min Calcium	1.0%
Min Crude Fat	2.5%	Min Available Phos	0.5%
Max Crude Fibre	6.0%	Min Added Copper	8.0mg
Max Added Salt	0.5%	Min Added Selenium	0.3mg
		Bovatec® 20CC (Lasalocid Sodium)	80mg



WHAT'S IN IT?

INGREDIENTS SELECTED FROM THE FOLLOWING: Cereal Grains and their By-products, Legumes and their By-products, Vegetable Protein Meals, Fats and Oils, Calcium Carbonate, Mono-Di Calcium Phosphate, Ridley Vitamin and Mineral Premix, Essential Amino Acids and Enzymes, & the Anticoccidial - Bovatec® 20CC.



PROUDLY MADE IN AUSTRALIA
USING INGREDIENTS SOURCED
FROM AUSTRALIAN FARMERS.

EXPECTED DAILY FEED INTAKE

50g per bird per day.

FEEDING GUIDE

- Ensure birds have continuous access to feed - do not restrict consumption.
- Self feeders designed for poultry are a good way to ensure feed is continuously available.
- Hatchlings can be encouraged to start feeding by scattering feed on newspaper placed around the feeders.
- Ensure that birds have access to clean, fresh water at all times.
- Follow any other feeding instructions given to you by your livestock provider.

This feed has been formulated specifically for animals as described on the package, or the label attached to the package, and must not be used for other species of animals.
ANIMAL FOODSTUFF. UNSUITABLE FOR HUMAN CONSUMPTION.

This product does not contain Restricted Animal Material.

CONTAINS MEDICATION: Contains 80mg/kg Lasalocid Sodium (as Bovatec® 20CC Lasalocid Sodium Premix) as an aid in the prevention of coccidiosis caused by *Eimeria sp* in layer replacement pullets. Bovatec® 20CC Lasalocid Sodium Premix is a registered trademark of Zoetis Inc.

WARNING: Bovatec® 20CC Lasalocid Sodium Premix is not registered for use in laying hens (chickens or turkeys). DO NOT feed to horses or other equids. DO NOT feed to dogs.

WITHHOLDING PERIOD: Meat - nil. Broilers and replacement Pullets: DO NOT USE in broilers or replacement pullets less than 3 days before slaughter for human consumption. Eggs - DO NOT use in laying hens within 14 days of laying where eggs or egg products are to be used for human consumption. In this instance introduce a Barastoc layer feed 2 weeks prior to the point of lay.

The information provided in this specification is exclusive to the product identified and is accurate at the time of writing. The manufacturer accepts no responsibility if this feed is used in quantities, conditions or in any way other than in accordance with the manufacturer's recommendations.

STORAGE INSTRUCTIONS

For optimum quality and shelf life, product should be stored below 30°C and off the floor in a cool, dry, well ventilated area protected from direct sunlight.



SCAN THE QR CODE
to visit the Barastoc Chook Book,
a comprehensive guide on rearing,
feeding and keeping poultry.



we put the chicken before the egg