BARASTOC

ESSENTIAL FEEDS

CHICK STARTER

COMPLETE & BALANCED CRUMBLE





WHAT WILL IT PROVIDE MY CHICKEN?

Each pellet is a concentrate of high quality ingredients specifically formulated to support the growth and health of chicks during the first few weeks post hatching.

Contains an anticoccidial as an aid in the prevention of coccidiosis*, an intestinal infection caused by the Eimeria parasite which young birds are vulnerable to.

Presented as a small crumble which is suitable for small beaks.

NUTRIENT LEVELS

TYPICAL ANALYSIS AS FED PER KG

| NUTRIENT LEVELS | | | |
|-------------------|-------|--------------------|-------|
| Min Crude Protein | 19.5% | Min Available Phos | 0.5% |
| Min Crude Fat | 2.5% | Min Added Copper | 8.0mg |
| Max Crude Fibre | 6.0% | Min Added Selenium | 0.3mg |
| Max Added Salt | 0.5% | Lasalocid Sodium | 80mg |
| Min Calcium | 1.0% | | |





WHAT'S IN IT?

INGREDIENTS SELECTED FROM THE FOLLOWING: Cereal Grains and their By-products, Legumes and their By-products, Vegetable Protein Meals, Fats and Oils, Calcium Carbonate, Mono-Di Calcium Phosphate, Salt, Ridley Vitamin and Mineral Premix, Essential Amino Acids and Enzymes, Anticoccidial - Bovatec* 20CC.



EXPECTED ADULT DAILY FEED INTAKE

50g per bird per day.

FEEDING GUIDE

Ensure birds have continuous access to feed - do not restrict consumption. Expected chick intakes are 50g/bird/day.

Self feeders designed for poultry are a good way to ensure feed is continuously available.

Ensure birds have access to clean, fresh water at all times.

Hatchlings can be encouraged to start feeding by scattering feed on newspaper placed around feeders.

Follow any other feeding instructions given to you by your livestock provider.

The manufacturer accepts no responsibility if this feed is used in quantities, conditions or in any way other than in accordance with the manufacturer's recommendations.

The manufacturer's recommendations for the use of this product may be obtained by contacting the Ridley Sales & Support Office on 1300 666 657, or the registered office of the company.

STORAGE REQUIREMENTS: For optimum quality and shelf life product should be stored below 30°C and off the floor in a cool, dry, well ventilated area protected from direct sunlight.

WARNING: THIS FEED HAS BEEN FORMULATED SPECIFICALLY FOR ANIMALS AS DESCRIBED ON THE PACKAGE, OR THE LABEL ATTACHED TO THE PACKAGE,

AND MUST NOT BE USED FOR OTHER SPECIES OF ANIMALS. UNSUITABLE FOR HUMAN CONSUMPTION.

CONTAINS MEDICATION: Contains 80mg/kg Lasalocid Sodium (as Bovatec* 20CC Lasalocid Sodium Premix) as an aid in the prevention of coccidiosis caused by Eimeria spp in layer replacement pullets.

WARNING: ONLY for broilers, replacement pullets and turkeys. DO NOT FEED to hens producing fertile eggs or to game birds other than turkeys. DO NOT feed to horses or other equids.

WITHHOLDING PERIOD: MEAT – DO NOT USE less than 1 day before slaughter for human consumption. EGGS: DO NOT USE in laying birds within 14 days of laying where eggs or egg products are to be used for human consumption or processing. TURKEYS: MEAT: Zero (0) days.



^{*}For more information about this condition refer to the Barastoc Chook Book.