

BARASTOC

TABLE-BIRD FEEDS

TURKEY & MEAT CHICKEN GROWER

COMPLETE & BALANCED GROWING CRUMBLE



WHAT WILL IT PROVIDE MY POULTRY?

Nutritionally balanced feed specifically formulated to help turkey poults and meat chickens develop into strong, healthy adults.

Contains an anti-coccidial as an aid in the prevention of coccidiosis, an intestinal infection caused by the Eimeria parasite which young birds are vulnerable to.

Presented as a small crumble which is suitable for small beaks.

NUTRIENT LEVELS

TYPICAL ANALYSIS AS FED PER KG

NUTRIENT LEVELS		TYPICAL ANALYSIS AS FED PER KG	
Min Crude Protein	20.0%	Min Available Phos	0.47%
Min Crude Fat	2.8%	Min Added Copper	8.0mg
Max Crude Fibre	6.0%	Min Added Selenium	0.3mg
Max Added Salt	0.2%	Lasalocid Sodium	100mg
Min Calcium	0.95%		



WHAT'S IN IT?

INGREDIENTS SELECTED FROM THE FOLLOWING: Cereal Grains and their By-products, Legumes and their By-products, Vegetable Protein Meals, Fats and Oils, Mono-Di Calcium Phosphate, Calcium Carbonate, Essential Amino Acids and Enzymes, Ridley Mineral and Vitamin Premix, Anticoccidial - Bovatec® 20CC.



MADE IN AUSTRALIA BY A PROUDLY AUSTRALIAN OWNED COMPANY.

EXPECTED DAILY FEED INTAKE

100 - 105g per chicken poult per day. 400g per turkey poult per day.

FEEDING GUIDE

Ensure birds have continuous access to feed - do not restrict consumption. Self-feeders designed for poultry are a good way to ensure feed is continuously available.

Ensure that birds have access to clean, fresh water at all times.

Follow any other feeding instructions given to you by your livestock provider.

The manufacturer accepts no responsibility if this feed is used in quantities, conditions or in any way other than in accordance with the manufacturer's recommendations. The manufacturer's recommendations for the use of this product may be obtained by contacting the Ridley Sales and Support Office on 1300 666 657 or the Head Office of the company.

STORAGE REQUIREMENTS:

For optimum quality and shelf life, product should be stored below 30°C and off the floor in a cool, dry, well ventilated area protected from direct sunlight.

WARNING: THIS FEED HAS BEEN FORMULATED SPECIFICALLY FOR ANIMALS AS DESCRIBED ON THE PACKAGE, OR THE LABEL ATTACHED TO THE PACKAGE, AND MUST NOT BE USED FOR OTHER SPECIES OF ANIMALS. ANIMAL FOODSTUFF. UNSUITABLE FOR HUMAN CONSUMPTION.

CONTAINS MEDICATION: Contains 100mg/kg Lasalocid Sodium (as Bovatec® 20CC Lasalocid Sodium Premix) as an aid in the prevention of coccidiosis caused by Eimeria sp in chicken broilers and turkey poults.

WARNING: Bovatec® 20CC Lasalocid Sodium Premix is not registered for use in laying hens (chickens or turkeys). DO NOT feed to horses or other equids. DO NOT feed to dogs.

WITHHOLDING PERIOD: Meat - nil. Broilers and replacement Pullets: DO NOT use in broilers or replacement pullets less than 3 days before slaughter for human consumption. Eggs - DO NOT use in laying hens within 14 days of laying where eggs or egg products are to be used for human consumption. In this instance introduce a Barastoc layer feed 2 weeks prior to the point of lay.



For more information on any of our products, call: 1300 666 657
visit: www.barastocpoultry.com.au • email: enquiries@ridley.com.au

we put the chicken before the egg